

Supplementary Appendix

Supplementary Appendix Table of Contents

LIBERTY 1 Principal Investigators and Study Sites	3
LIBERTY 2 Principal Investigators and Study Sites	9
Supplementary Methods	16
Selection of Study Population	16
Table S1. Eligibility Criteria.	16
Study Objectives and End Points	20
Table S2. Study Objectives and End Points.	20
Mixed-Effects Model for Imputing Missing Menstrual Blood Loss Volume Data	24
Derivation of Responder Status at Week 24/End of Treatment and Missing Data Handling Rules	26
Table S3. Derivation of Responder Status at Week 24/End of Treatment and Missing Data Handling Rules – for Primary Analysis.	28
Key Secondary Efficacy End Points with Alpha-Protection	29
Figure S1. Mixed Sequence Testing Procedure for Primary and Key Secondary End Points in LIBERTY 1.	30
Figure S2. Mixed Sequence Testing Procedure for Primary and Key Secondary End Points in LIBERTY 2.	31
Patient-Reported Outcomes	34
Supplementary Figures	37
Figure S3. Study Design.	37
Figure S4. Patient Disposition.	38
Figure S5. Subgroup Analyses for Primary End Point	39
Figure S6. Secondary Efficacy End Points: A) Reduction in Menstrual Blood Loss, B) Relief of Pain Associated With Uterine Fibroids, and C) Reduction in Bleeding and Pelvic Discomfort.	40
Supplementary Tables	41
Table S4. Additional Baseline Characteristics.	41
Table S5. Other Secondary End Points.	42
Table S6. Serious Adverse Events.	43
Table S7. Percent Change from Baseline in Bone Mineral Density at Lumbar Spine and Total Hip in LIBERTY 1 and LIBERTY 2.	44
Table S8. Summary of Patients With Liver Function Tests Meeting Predefined Limits of Change.	45
Table S9. Summary of Change from Baseline in Serum Lipids.	46
References	48

LIBERTY 1 Principal Investigators and Study Sites

Site	Investigator Name	Facility Name	Address
United States			
1000	Samuel Alexander, MD	Southern Clinical Research Associates	4720 South I-10 Service Rd Suite 201 Metairie, LA 70001
1001	Mark Firestone, MD	South Florida Medical Research	21150 Biscayne Blvd Suite 300 Aventura, FL 33180
1002	Frances Fisk, MD	Southwest Clinical Research	4901 Lang Ave NE Suite 203 Albuquerque, NM 87109
1003	Dana Shipp, MD	Medical Center for Clinical Research	9040 Friars Road Suite 540 San Diego, CA 92108
1004	Andrea Lukes, MD	Carolina Women's Research and Wellness Center	249 E NC Highway 54 Suite 330 Durham, NC 27713
1005	R Garn Mabey, MD	Rex Garn Mabey, Jr, MD Chartered	2881 N Tenaya Way Las Vegas, NV 89128
1006	Marina Raikhel, MD	Torrance Clinical Research Institute, Inc.	25043 Narboone Ave Lomita, CA 90717
1007	Steven Sussman, MD	Lawrence OB/GYN Clinical Research, LLC	123 Franklin Corner Rd Suite 214 Lawrenceville, NJ 08648
1008	Stephen Gene Swanson, MD	Women's Clinic of Lincoln, PC	220 Lyncrest Dr Lincoln, NE 68510
1009	Arthur Waldbaum, MD	Downton Women's Health Care	3773 Cherry Creek Dr N # 685 Denver, CO 80209
1010	Ronald Ackerman, MD	Comprehensive Clinical Trials, LLC	603 Village Blvd Suite 301 Memphis, TN 38119
1026	Samuel Simha, MD	Research Memphis Associates, LLC	1028 Cresthaven Rd Suite 101 Memphis, TN 38119
1027	Pouru Bhiwandi, MD	Wake Research Associates, LLC	3100 Duraleigh Rd Suite 304 Raleigh, NC 27612
1028	Sandra Hurtado, MD	The Woman's Hospital of Texas Clinical Research Center	7400 Fannin St Suite 1280 Houston, TX 77054
1029	Stuart Weprin, MD	HWC Women's Research Center, LLC	20 West Wenger Rd Englewood, OH 45322
1030	Albert Tydings, MD	Clinical Trials Management, LLC	7043 Hwy 190 E Service Rd Suite C Covington, LA 70433
1031	Edward Zbella, MD	Women's Medical Research Group, LLC	2454 McMullen Booth Rd Suite 609 Clearwater, FL 33759
1032	Carlito Arrogante, MD	South Florida Clinical Trials	7100 W 20th Avenue Suite 701 Hialeah, FL 33016

Site	Investigator Name	Facility Name	Address
1033	Kevin Fleishman, MD	Clinical Physiology Associates	13670 Metropolis Ave Suite 105 Fort Myers, FL 33912
1034	Patricia Thompson, MD	Fellows Research Alliance, INC	11 Arley Way Suite 202 Bluffton, SC 29910
1035	Ivonne Reynolds, DO	South Florida Clinical Research Institute	2964 N State Road 7 Suite 200 Margate, FL 33063
1044	Beatrice Hecker, MD	Miami Dade Medical Research Institute, LLC	8955 SW 87th Ct Suite 112 Miami, FL 33176
1045	Andrew Kaunitz, MD	University of Florida College of Medicine-Jacksonville, Dept. of Obstetrics and Gynecology	4549 Emerson Street Bldg 2, Suite 201 Jacksonville, FL 32207
1046	Guillermo Lievano, DO	Genoma Research Group	2720 SW 97th Ave Suite 102 and 103 Miami, FL 33165
1047	Kathryn Moore, MD	Vista Clinical Research	700 Gervais St, Suite 300 Columbia, SC 29201
1049	Nasiruddin Rana, MD	Affinity Clinical Research Institute	2425 W 22nd St Suite 216 Oakbrook, IL 60523
1050	Sharifa Fazili, MD	Upstate Clinical Research Associates, LLC	8201 Main Street Suite 1 Williamsville, NY 14221
1052	Winifred Soufi, MD	Agile Clinical Research Trials	980 Johnson Ferry Rd Suite 720 Atlanta, GA 30342
1069	Danilo Herrera, MD	Coastal Clinical Research	100 Memorial Hospital Dr Annex Building, Suite 3-B Mobile, AL 36608
1071	Sandra Nieto, MD	Futura Research Inc.	11514 Rosecrans Ave Norwalk, CA 90650
1072	Rovena Reagan, MD	Women's Health Care Research Corp.	3750 Convoy St Suite 311 San Diego, CA 92111
1073	Marlin Schul, MD	Lafayette Clinical Research	3920 St. Francis Way Suite 107 Lafayette, IN 47905
1074	Sherry Soefje, MD	Excell Research, Inc	3998 Vista Way Suite 100 Oceanside, CA 92056
1075	John Whitfield, MD	Signature Gyn Services	1250 8th Ave Suite 540 Fort Worth, TX 76104
1102	James Roth, MD	Advanced Clinical Research/Old Farm Obstetrics& Gynecology	3970 South 700 East Suite 14 Salt Lake City, UT 84124
1103	David F. Archer, MD	Eastern Virginia Medical School	601 Colley Ave Rm 241 Norfolk, VA 23507

Site	Investigator Name	Facility Name	Address
1104	Ladynez Espinal, MD	Clinovation Research, LLC	2700 Glades Circle #142 Weston, FL 33327
1121	Kirk Brody, MD	Chattanooga Medical Research	2341 McCallie Avenue Chattanooga, TN 37404
1122	Hugo Perez, MD	Suncoast Clinical Research	3890 Tampa Road Suite 301 Palm Harbor, FL 34684
1123	Nelson Uzquiano, MD	UAG Innovation Women Research, LLC	7737 Southwest Freeway Suite 870 Houston, TX 77074
1125	Michael A. Wells, MD	SEC Clinical Research	822 South Three Notch St Suite B Andalusia, AL 36420
1127	John Cote, MD	CHI Health Research Center	7710 Mercy Road Suite 228 Omaha, NE 68124
1128	Parke Hedges, MD	Clinical Trials of Texas, Inc.,	7940 Floyd Curl Drive Suite 700 San Antonio, TX 78229
1129	Carolyn W. Quist, DO	Brownstone Clinical Trials	1425 8th Ave Suite 101 Fort Worth, TX 76104
1168	Anthony Adams, MD	Paramount Research Solutions/Apogee Women's Health	2565 Jolly Road Suite B College Park, GA 30349
1169	Kwabena Ayesu, MD	Omega Research Consultants, LLC	7912 Forest City Rd Suite 101 Orlando, FL 32810
1170	Eric Brown, MD	Medisense, Inc	2635 South Cobb Drive Smyrna, GA 30080
1173	Fred Newton, MD	Lynn Institute of the Ozarks	500 S University Ave Suite 500 Little Rock, AR 72205
1191	Elia Fanous, MD	Brownstone Clinical Trials	3501 N MacArthur Blvd Suite 500 Irving, TX 75062
1193	Robin Kroll, MD	Seattle Women's: Health, Research, and Gynecology	3216 NE 45th Place Suite 100 Seattle, WA 98105
1194	Gioi Smith-Nguyen, MD	Grossmont Center for Clinical Research	8851 Center Dr Suite 206 La Mesa, CA 91942
1195	Kurt Barnhart, MD	Department of Obstetrics and Gynecology, University of Pennsylvania School of Medicine	3701 Market Street 8th Floor Philadelphia, PA 19104
1197	Aparna Tamirisa, MD	Houston Center for Clinical Research	3519 Town Center Blvd Suite B 2nd Floor Sugar Land, TX 77479
1205	Todd B. Chappell, MD	Adams Patterson GYN/Wake Research Associates	6215 Humphreys Blvd Suite 301 Memphis, TN 38120

Site	Investigator Name	Facility Name	Address
1206	Vicki Schnell, MD	Center for Reproductive Medicine	1015 Medical Center Blvd Suite 2100 Webster, TX 77598
1209	Molly Katz, MD	Katz and Kade, Inc.	71 E Hollister St Cincinnati, OH 45219
1221	Mara Rabin (fka Epstein), MD	Wasatch Clinical Research	4001 South 700 East Suite 105 Salt Lake City, UT 84107
1222	Ronnie M Givens II, MD	Clinical Trials of South Carolina	2695 Elms Plantation Blvd Suite D Charleston, SC 29406
1223	Elliot E Cazes, MD	New Tampa Ob/GYN (Stedman Clinical Trials)	3212 Cove Bend Dr Tampa, FL 33613
1233	Julie Saranita, DO	South Lake Pain Institute (Clin-Edge - Chris Matteo/Christian Burns)	2440 Hooks St Clermont, FL 34711
1239	Kelli Braun, MD	Augusta University Medical Center	1521 Pope Avenue Augusta, GA 30912
1240	Celestino Castellon, MD	Direct Helpers Medical Center, Inc.	Hialeah, FL
1253	Keila Hoover, MD	Miami Clinical Research	7371 SW 24 th Street Augusta, GA 30912
Rest of World			
Brazil			
1099	Emerson de Oliveira	Centro Multidisciplinar de Estudos Clínicos Unidade Santo André	Avenida Gilda 106 Salas 82 e 83 Centro Santo André, Sao Paulo, Brazil 0910-510
1142	Luciano Gibran	Hospital Pérola Byington - Centro de Referência da Saúde da Mulher	Avenida Brigadeiro Luís Antônio, 683 6 andar - sala 600 Bela Vista São Paulo, Sao Paulo, Brazil 01317-000
1167	Carlos Isaia Filho	Unidade de Pesquisa Clínica - Centro de Medicina Reprodutiva Dr. Carlos Isaia Filho	Hilario Ribeiro 202/304 Porto Alegre, Rio Grande Do Sul, Brazil 90510-040
Italy			
1091	Fiorenzo, De Cicco Nardone	Fondazione Policlinico Universitario Agostino Gemelli	Clinica Ostetrica e Ginecologica Largo Agostino Gemelli 8 Roma, Roma, Italy 00168
1093	Vincenzo De Leo, MD	Azienda Ospedaliera Universitaria Senese	Unità Operativa Complessa Viale Mario Bracci 16 Siena, Siena, Italy 53100
1101	Maria Grazia Porpora	Universita degli Studi di Roma "La Sapienza" - Umberto I Policlinico di Roma	Ostetricia e Ginecologia Dipartimento Scienze Ginecologico-Ostetriche Viale Regina Elena 324 Roma, Roma, Italy 00161

Site	Investigator Name	Facility Name	Address
3094	R. Venturella	Azienda Ospedaliera "Pugliese Ciaccio" di Catanzaro	Dipartimento Integrazione e Tutela Maternità ed Infanzia Ostetricia e Ginecologia Universitaria Via Papa Pio X, 83 Catanzaro, Italy 88100
3123	M. Bau	Azienda Ospedaliero Universitaria "Citta della Salute e dell Scienza" di Torino – Ospedale Sant Anna	SC Ginocologia e Ostetricia 3 Corso Spezia, 60 Torino, Italy 10126
3125	F. Petraglia	Azienda Ospedaliero Universitaria Careggi	DAI Meterno – Infantile SOD Ostetricia e Ginecologia Largo Brambilla, 3 Firenze, Italy 50134
Poland			
1151	Tomasz Rechberger, Prof	Niepubliczny Zakład Opieki Zdrowotnej NOVITA Specjalistyczne Gabinety Lekarskie	ulica Jana Sawy 1B/5,6 Lublin, Lubelskie, Poland 20-632
1152	Zoulikha Jabiry-Zieniewicz	NZOZ Zieniewicz Medical	ulica Opaczewska 43/18 Warszawa, Mazowieckie, Poland 02-0201
1156	Krzysztof Wilk	Wielospecjalistyczna Poradnia Lekarska Synapsis	ulica Czerwinskiego 8 Katowice, Slaskie, Poland 40-123
1159	Slawomir Jedrzejczyk	Niepubliczny Zakład Opieki Zdrowotnej JORDAN Slawomir Jedrzejczyk	ulica Zielona 29 Łódź, Lodzkie, Poland 90-602
3149	K. Oronowicz	Twoja Przychodnia – Szczecińskie Centrum Medyczne	Słowackiego 19, Szczecin, Poland 71-434
South Africa			
1160	Sagaren Naidu	Dr S Naidu O and G Clinical Trials	Nordica Durban Fertility Clinic Netcare Kingsway Hospital Suite B3 607 Andrew Zondo Road Durban, KwaZulu-Natal, South Africa 4126
1236	Elmarie Wypkema	Life Wilgeheuwel Hospital	Amplifier Road Radiokop Ext 13 Block A Suite 31 Roodepoort, Gauteng, South Africa 1724
1238	Daniel Malan	Phoenix Pharma	2 Eastbourne Road Mount Croix Port Elizabeth, Eastern Cape, South Africa 6001

Site	Investigator Name	Facility Name	Address
United Kingdom			
1188	Nitish Narvekar	King's College Hospital NHS Foundation Trust	Assisted Conception Unit First Floor, Unit 6, KCH Business Park Coldharbour Lane London, England, United Kingdom SE5 9NY

LIBERTY 2 Principal Investigators and Study Sites

Site	Investigator Name	Facility Name	Address
United States			
1011	Jeffrey B Baker, MD	Clinical Research Prime (CR Prime)	187 East 13th Street Idaho Falls, ID 83404
1013	Robert E Deck Jr., MD	Clinical Trials Management, LLC	3801 Houma Blvd. Suite 200 Metairie, LA 70006
1014	Robert Smith Jr., MD	Suncoast Clinical Research	5604 Gulf Dr. New Port Richey, FL 34652
1015	Cynthia Goldberg, MD	Visions Clinical Research - Tucson	5225 E. Knight Drive Suite #101 Tucson, AZ 85712
1016	Ernesto Gomez, MD	Mesa Obstetricians and Gynecologists	7233 E. Baseline Rd. Suite 101 Mesa, AZ 85209
1017	Debra Walland,, MD	Fellows Research Alliance, Inc.	1 Oglethorpe Professional Blvd. Suite 204 Savannah, GA 31406
1019	Gwang-Yi Hwang, MD	Physicians' Research Options, Red Rocks Ob/Gyn	255 Union Blvd. Suite 200 Lakewood, CO 80228
1020	Edmond Pack, MD	Office of Edmond E. Pack	8285 W Arby Ave Suite 280 Las Vegas, NV 89113
1021	Khadra Osman, MD	KO Clinical Research, LLC	1625 SE 3rd Ave Suite 400 Ft. Lauderdale, FL 33316
1022	Phyllis Gee, MD	Willowbend Health & Wellness Associates	4401 Coit Rd. Suite 205 Frisco, TX 75035
1023	Michael Swor, MD	Physician Care Clinical Research, LLC	1617 South Tuttle Avenue Suite 1A Sarasota, FL 34239
1024	Alfred Poindexter, III, MD	Advances in Health	7515 South Main Suite 360 Houston, TX 77030
1040	Charles Newlin, MD	DCOL Center for Clinical Research	707 Hollybrook Dr. Suite 501 Longview, TX 75605
1041	Ronald Surowitz, DO	Health Awareness, Inc.	1094 Military Trail 2nd Floor Jupiter, FL 33458
1043	James Simon, MD	James A. Simon, MD, PC	1850 M Street NW Suite 450 Washington, DC 20036
1053	Vaughn Whittaker, MD	New York Clinical Trials - Manhattan (NYCT, A Member of the Alliance, Inc.)	59 East 54th Street 9th Floor, Suite 93 New York, NY 10022

Site	Investigator Name	Facility Name	Address
1054	Juana Cuevas, MD	New York Clinical Trials- Brooklyn (NYCT, A Member of the Alliance, Inc.)	50 Court Street 10th Floor, Room 1002 Brooklyn, NY 11201
1056	Brian MacGillivray, MD	Stone Oak, LLC dba Discovery Clinical Trials	19054 Stone Oak Pkwy Suite 102 San Antonio, TX 78258
1057	Charles E Miller, MD	The Advanced Gynecologic Surgery Institute	120 Osler Drive Suite 100 Naperville, IL 60540
1058	Andrew Wagner, MD	Saginaw Valley Medical Research Group, LLC	5400 Mackinaw Road Suite 6100 Saginaw, MI 48604
1076	Steven E. Chavoustie, MD	Healthcare Clinical Data, Inc.	1065 NE 125th Street Suite 102 North Miami, FL 33161
1077	Lydie L. Hazan, MD	Axis Clinical Trials (ACTCA, A Member of the Alliance, Inc.)	5800 Wilshire Blvd. Los Angeles, CA 90036
1078	Jody Jones, MD	ClinSite, LLC	49650 Cherry Hill Road Suite 220 Canton, MI 48187
1079	Clifton Nicholson-Uhl, MD	Praetorian Pharmaceutical Research, LLC	4700 Wichers Drive Suite 200 Marrero, LA 70072
1080	Nancy Phillips, MD	Rutgers-Robert Wood Johnson Medical School	125 Paterson Street New Brunswick, NJ 08901
1081	Dominique Smith, MD	Soapstone Center for Clinical Research	4201 Rainbow Drive Decatur, GA 30034
1082	Franklin Morgan, Jr., MD	Tidewater Physicians for Women	844 Kempsville Road Suite 208 Norfolk, VA 23502
1083	Ronald Hardy, MD	North Spokane Women's Health Research	235 E. Rowan Suite 206 Spokane, WA 99207
1084	Kord Strebhel, MD	Nevada Obstetrical Charity Clinic	1950 Pinto Lane Las Vegas, NV 89106
1105	Jay Cohen, MD	All Women's Healthcare of West Broward, Inc.	140 SW 84th Street Suite D Plantation, FL 33324
1106	Martin Kabongo, MD	Precision Research Institute	292 Euclid Ave. Suite 115 San Diego, CA 92114
1108	Antoinette Pragalos, MD	CTI Clinical Research Center	1775 Lexington Avenue Suite 200 Cincinnati, OH 45212
1133	Delbert "Alan" Johns, MD	Texas Health Care, PLLC	1250 8th Avenue Suite 330 Fort Worth, TX 76104
1134	Marigene Salazar Sharma, MD	Albuquerque Clinical Trials, Inc.	711 Encino Place, NE Albuquerque, NM 87102
1135	Danielle A. "Dee-Dee" Shiller, DO	Pharmasite Research, Inc.	1314 Bedford Avenue Suite 205 Baltimore, MD 21208

Site	Investigator Name	Facility Name	Address
1161	Jay Berman, MD	C.S. Mott Center- Wayne State University	275 E. Hancock 1st Floor Detroit, MI 48201
1162	Gordon B. Clark, MD	GTC Research	21624 Midland Drive Shawnee, KS 66218
1163	Suzanne L. Fussell, MD	Long Beach Clinical Trial Services, Inc.	2403 Atlantic Avenue Long Beach, CA 90806
1164	Milroy J. Samuel, MD	Complete Healthcare for Women	5888 Cleveland Avenue Columbus, OH 43231
1165	Arthur Schatz, MD	Ideal Clinical Research	2627 NE 203rd Street Suite 100 Aventura, FL 33180
1176	Robert Littleton, MD	Unified Women's Clinical Research-Raleigh	4414 Lake Boone Trail Suite 205 Raleigh, NC 27607
1177	Silpa Senchani, MD	ONCOVA Clinical Research, Inc.	3129 Innovation Drive St. Cloud, FL 34769
1178	Hugh A. Wilkinson, MD	Axcess Medical Research	13005 Southern Blvd. Bldg #2, Suite 214 Loxahatchee, FL 33470
1198	Clark Griffith, MD	DCT - HCWC, LLC, dba Discovery Clinical Trials	8160 Walnut Hill Lane Margot Perot Bldg., Suite 116 Dallas, TX 75231
1199	Amber G. Hatch, MD	Unified Women's Clinical Research d/b/a Lyndhurst Clinical Research	111 Hanestown Court Suite 151 Winston Salem, NC 27103
1201	David DeAtkine, Jr., MD	Central Research Associates, Inc.	2660 10th Avenue South Bldg. 1, Suite 735 Birmingham, AL 35205
1202	Juan P. Frias, MD	National Research Institute	2010 Wilshire Blvd, Suite 302 Los Angeles, CA 90057
1213	James P. Rice, MD	Multicare Women's Healthcare	17700 SE 272nd Street Suite 105 Covington, WA 98042
1215	Yamirka Duardo Guerra, MD	LCC Medical Research Institute, LLC	1150 NW 72nd Ave. Suite 620 Miami, FL 33126
1228	Jared Robins, MD	Northwestern University Division of Reproductive Endocrinology and Infertility Arkes Pavilion	676 N. St. Clair Street Suite 1845 Chicago, IL 60611
1232	David Montaldi, DO	One Health Research Clinic, Inc.	5880 Live Oak Pkwy Suite 160 Norcross, GA 34996
1249	Issam Daya, MD	Continental Clinical Solutions, LLC	7801 York Road Suite 203 Towson, MD 21204
1251	Bradley Block, MD	Oviedo Medical Research	2441 West State Road 426 Suite 2011 Oviedo, FL 32765

Site	Investigator Name	Facility Name	Address
1257	Brian Siu, MD	Marvel Clinical Research	17822 Beach Blvd Suite 325 Huntington Beach, CA 92647
1258	Gregory P. Moore, II, MD	Clinical Research Partners, LLC	110 North Robinson Street Suite 200 Richmond, VA 23225
1260	Hessam Aazami, MD	Hope Clinical Research	22030 Sherman Way Suite 101 Canoga Park, CA 91303
1262	Parvin Syal, MD	National Research Institute	14418 Chase Street Suite 200 Panorama City, CA 91402
1266	Ronald Ackerman, MD	Comprehensive Clinical Trials, LLC	603 Village Blvd. Suite 301 West Palm Beach, FL 33409
1267	Keila Hoover, MD	Miami Clinical Research	7371 SW 24th Street Miami, FL 33155
1268	Sandra Hurtado, MD	The Woman's Hospital of Texas Clinical Research Center	7400 Fannin St Suite 1280 Houston, TX 77054
1270	Andrea Lukes, MD	Carolina Women's Research and Wellness Center	249 E NC Highway 54. Suite 330 Durham, NC 27713
1273	Nasiruddin Rana, MD	Affinity Clinical Research Institute	2425 W 22nd St Suite 209 Oakbrook, IL 60523
3006	Godson Oguchi, MD	Midland Florida Clinical Research Group	665 Peachwood Drive Deland, FL 32720
3007	Stephen Blank, MD	Mount Vernon Clinical Research, LLC/	755 Mount Vernon Highway, NE Suite 200 Sandy Springs, GA 30328
3013	Marquita Anderson, MD	Gadolin Research LLC	2965 Harrison Avenue Suite 313 Beaumont, TX 77702
3175	Anthony Imudia, MD	USF Health, Department of Obstetrics & Gynecology	2 Tampa General Circle STC 4th Floor Tampa, FL 33606
3206	Duane Anderson, MD	Excel Clinical Research	3059 South Maryland Parkway Suite 202 Las Vegas, NV 89109
Rest of World			
Belgium			
1150	Jean-Luc Squifflet	Cliniques Universitaires Saint-Luc	Avenue Hippocrate 10 Brussels, Brussels, Belgium 1200
1157	Wim Decler	Algemeen Ziekenhuis Jan Palfijn Gent	Algemeen Ziekenhuis Jan Palfijn Gent IVF centre Gent, Oost-Vlaanderen, Belgium 9000

Site	Investigator Name	Facility Name	Address
1158	Stefan Cosyns	Universitair Ziekenhuis Brussel	Kliniekhofd Gynecologie – Oncologie Laarbeeklaan 101 Jette, Brussels, Belgium 1090
1226	Michaël De Brucker	Centre Hospitalier Universitaire de Tivoli	Avenue Max Buset 34 La Louvière, Hainaut, Belgium 7100
Brazil			
1061	Sonia Debbio, MD	Centro Paulista de Investigação Clínica	Rua Moreira e Costa 342 Ipiranga São Paulo, Sao Paulo, Brazil 04266-010
1063	Maria Celeste Wender, MD, PhD, ScD	Hospital de Clínicas de Porto Alegre	Rua Ramiro Barcelos 2350 Porto Alegre, Sao Paulo, Brazil 90035-903
1095	Carlos Isaia Filho	Unidade de Pesquisa Clínica - Centro de Medicina Reprodutiva Dr. Carlos Isaia Filho	Hilario Ribeiro 202/304 Porto Alegre, Rio Grande Do Sul, Brazil 90510-040
1096	Daniel Dias	Unidade de Pesquisa Clínica da Faculdade de Medicina de Botucatu	Rua Prof. Dr. Armando Alves s/n UNESP Câmpus de Botucatu Botucatu, Sao Paulo, Brazil 18618-686
Chile			
1112	Claudio Andrés Villarroel Quintana	Institute for Mother and Child Research (IDIMI), faculty of Medicine, University of Chile Hospital Clinico San Borja Arriaran	Avenida Santa Rosa 1234 Santiago, Santiago, Chile 8360160
1116	Guillermo Galan Chiappa	Centro de Capacitación e Investigaciones Clínicas	Guardia Vieja Oficina 207 Providencia, Santiago, Chile 11
1148	Cristian Jesam Gaete	Instituto Chileno de Medicina Reprodutiva	Avenida Santa Rosa 1234 Santiago, Santiago, Chile 8360160
1246			
Czech Republic			
1065	Aleš Skrivánek	G-CENTRUM	Horní náměstí 285/8 Olomouc, Czech Republic 772 00
1097	Olga Hlaváčková	Centrum Gynekologické Rehabilitace	Zeyerova 2442/19 Písek, Jihočeský Kraj, Czech Republic 39701
1115	Aleš Roztocil	Nemocnice Jihlava	Vrchlického 59 Jihlava, Kraj Vysocina, Czech Republic 586 33
1146	Petr Sák, MD, PhD	MUDr. Petr Sák	Senovážné náměstí 248/2 České Budejovice, Czech Republic 370 01
Hungary			
1059	János Zatik	Szent Anna Nőgyógyászati, Szülészeti és Ultrahang Magánrendelő	Szent Anna utca 48 Debrecen, Hajdu-Bihar, Hungary 4024

Site	Investigator Name	Facility Name	Address
1064	Károly Pap, MD	Szabolcs-Szatmár-Bereg Megyei Kórházak és Egyetemi Oktatókórház	Szent István utca 68 Nyíregyháza, Szabolcs-Szatmar-Bereg, Hungary 4400
1068	József Bódis, MD	Pécsi Tudományegyetem Klinikai Központ	Édesanyák útja 17 Pécs, Baranya, Hungary 7624
1145	Tamás Nyirády	Mediroyal	Fráter György utca 4 Kecskemét, Bacs-Kiskun, Hungary 6000
1155	Ferenc Racz	Csongrád Megyei Dr. Bugyi István Kórház	Sima Ferenc utca 44-58 Szuleszeti es Nogyogyaszati Osztaly Szentes, Csongrad, Hungary 6600
3038	Márta Rákos	Synexus Magyarország Egészségügyi Szolgáltató Kft. - Affiliated Site Gyula	Nürnbergi utca 1/b, Gyula, Hungary 5700
3136	József Nemes	Synexus Magyarország	Miklós utca 5-13, Debrecen, Hungary 4025
Poland			
1085	Jacek Szamatowicz	ARTEMIDA Centrum Ginekologii Endokrynologii i Medycyny Rozrodu	Ul. Wlokiennicza 9b/u Bialystok, Podlaskie, Poland 15-464
1087	Krzysztof Szymanowski	Examen	ulica Barwicka 14 A Poznan, Wielkopolskie, Poland 60-192
1088	Janusz Tomaszewski, MD, PhD	Prywatna Klinika Polozniczo-Ginekologiczna	Parkowa 6 Bialystok, Podlaskie, Poland 15-224
1110	Tomasz Lozinski	Szpital Specjalistyczny Pro-Familia	ulica Witolda 6B Rzeszów, Podkarpackie, Poland 35-302
1244	Krzysztof Lukaszuk	Kliniki i Laboratoria Medyczne INVICTA	ul. Rajska 10 Gdansk, Pomorskie, Poland 80-850
South Africa			
1120	Tobias De Villiers	Clintrials Projects cc	1 Rothschild Boulevard 1st Floor Room 118 Mediclinic Panorama Cape Town, Western Cape, South Africa 7500
1138	Victor Hulme	Life Vincent Pallotti Hospital	Park Road Pinelands Cape Town, Western Cape, South Africa 7405
1147	Nyda Fourie	Iatros International	Quantum Building Middle Block - Ground Floor 1 Third Avenue, Westdene Bloemfontein, Free State, South Africa 9301

Site	Investigator Name	Facility Name	Address
1149	Lynette Reynders	Office Of Lynette Reynders, MB CHB	119 Cantonment Road Lyttelton Centurion, Gauteng, South Africa 0157

Supplementary Methods

Selection of Study Population

Table S1. Eligibility Criteria.

Inclusion Criteria
<p>A woman was eligible for randomization and enrollment in this study only if all of the following inclusion criteria applied and were met at the time of the baseline Day 1 visit, unless otherwise specified.</p>
<ol style="list-style-type: none"> 1. Had voluntarily signed and dated the informed consent form prior to initiation of any screening or study-specific procedures; 2. Was a premenopausal female 18 to 50 years of age (inclusive) on the day of signing and dating the informed consent form; 3. Had regularly-occurring menstrual periods of ≤ 14 days duration with a cycle of 21 to 38 days from the start of one menstrual period until the start of the next, by patient history for at least three months prior to the screening 1 visit; 4. Had a diagnosis of uterine fibroids that was confirmed by a transvaginal ultrasound performed during the screening period; at least one uterine fibroid had to be verified by a central reader to meet at least one of the following criteria: <ol style="list-style-type: none"> a. Subserosal, intramural, or $<50\%$ intracavitary submucosal fibroid with a diameter ≥ 2 cm (longest diameter), or b. Multiple small fibroids with a total uterine volume of ≥ 130 cm³
<p>Note: Once the transvaginal ultrasound was completed, a transabdominal ultrasound could have been performed if the uterus could not be adequately imaged on transvaginal ultrasound (i.e., because of enlarged size).</p>
<p>Note: Saline or gel contrast was not required but could have been performed to demonstrate fibroids that met the criterion for inclusion if these were not adequately visualized with transvaginal ultrasound alone.</p>
<ol style="list-style-type: none"> 5. Had heavy menstrual bleeding associated with uterine fibroids, as evidenced by a menstrual blood loss volume of ≥ 160 ml during 1 cycle or ≥ 80 ml per cycle for two menstrual cycles, as measured by the alkaline hematin method during the screening period; 6. Patient did not expect to undergo gynecological surgery or ablation procedures for uterine fibroids within the six months following enrollment; 7. Had a negative urine pregnancy test at the Screening 1, Screening 3, and baseline Day 1 visits; 8. Agreed to use contraception during the study and for 30 days following the last dose of study drug. Specifically agreed to use non-hormonal contraception consistently during the screening period and the randomized treatment period and either non-hormonal or oral contraceptives after return of menses following treatment discontinuation. However, the patient was not required to use specified non-hormonal contraception if the following applied: <ol style="list-style-type: none"> a. Had sexual partner(s) who were vasectomized at least six months prior to the screening period; b. Had a bilateral tubal occlusion (including ligation and blockage methods, such as Essure™), at least four months prior to the first screening visit (patients with Essure had to have prior confirmation of tubal occlusion by hysterosalpingogram and no evidence of “post-Essure syndrome,” in the investigator’s opinion); c. Was not sexually active with men; periodic sexual relationship(s) with men required the use

of non-hormonal contraception.

9. Had an adequate endometrial (aspiration) biopsy performed during the screening period, with results showing no clinically significant endometrial pathology (hyperplasia, polyp, or endometrial cancer).

Note: Patients for whom polyps were detected on biopsy but were either not evident on ultrasound or <2 cm were eligible;

10. If ≥ 39 years of age at the time of the baseline Day 1 visit, had a normal mammogram (Breast Imaging Reporting and Data System category 1 or 2 or equivalent) during the screening period or within six months prior to the screening period.

Exclusion Criteria

1. Had transvaginal and/or transabdominal ultrasound during the screening period demonstrating pathology other than uterine fibroids that could have been responsible for or contributed to the patient's heavy menstrual bleeding, such as uterine or cervical polyps ≥ 2 cm, large simple ovarian cyst >4.0 cm, endometrioma(s) >4.0 cm, or any other clinical significant gynecological disorder determined by the investigator to require further evaluation and/or treatment during the study;

Note: Saline or gel contrast was not routinely required. Use of such contrast was required only when the endometrium could not be evaluated or when there were ambiguous and potentially exclusionary findings on the transvaginal or transabdominal ultrasound (e.g., suspected intrauterine masses, equivocal endometrial findings, etc.);

2. Had known rapidly enlarging uterine fibroids, in the opinion of the investigator;
3. Had undergone myomectomy, ultrasound-guided laparoscopic radiofrequency ablation, or any other surgical procedure for fibroids, uterine artery embolization, magnetic resonance-guided focused ultrasound for fibroids, as well as endometrial ablation for abnormal uterine bleeding within six months prior to the screening 1 visit;
4. Had a weight that exceeded the weight limit of the dual-energy x-ray absorptiometry scanner or had a condition that precluded an adequate dual-energy x-ray absorptiometry measurement at the lumbar spine and proximal femur (e.g., bilateral hip replacement or spinal hardware in the lumbar spine);
5. Had a baseline bone mineral density z-score <-2.0 at the spine, total hip, or femoral neck;
6. Had a history of or currently had osteoporosis, or other metabolic disease, hyperparathyroidism, hyperprolactinemia, hyperthyroidism, anorexia nervosa, or low traumatic (from the standing position) or atraumatic fracture (toe, finger, skull, face and ankle fractures were allowed). Patients whose hyperparathyroidism or hyperthyroidism had been successfully treated or whose hyperprolactinemia had been successfully treated and/or who met bone mineral density eligibility criteria for the study were allowed;
7. Had a history of the use of bisphosphonates, calcitonin, calcitriol, ipriflavone, teriparatide, denosumab, or any medication other than calcium and vitamin D preparations to treat bone mineral density loss;
8. Anticipated use of systemic glucocorticoids at an oral prednisone-equivalent dose of more than 5 mg every other day during the study.

Note: topical, inhaled, intranasal, otic, ophthalmic, intraarticular, or intralesional subcutaneous glucocorticoids were permitted without restriction.

9. Gastrointestinal disorder affecting absorption or gastrointestinal motility;
10. Had any contraindication to treatment with estradiol and norethindrone acetate, including the following:
 - a. Known, suspected, or history of breast cancer;

- b. Known or suspected estrogen-dependent neoplasia;
 - c. Active deep vein thrombosis or pulmonary embolism, or history of these conditions prior to the baseline Day 1 visit;
 - d. History of or active arterial thromboembolic disease, including stroke and myocardial infarction;
 - e. Known anaphylactic reaction or angioedema or hypersensitivity to estradiol or norethindrone acetate;
 - f. Known protein C, protein S, or antithrombin deficiency, or other known thrombophilia disorders, including Factor V Leiden;
 - g. Migraine with aura;
 - h. History of porphyria;
11. Had jaundice or known current active liver disease from any cause, including hepatitis A, hepatitis B, or hepatitis C (hepatitis C virus [HCV] antibody positive, confirmed by HCV ribonucleic acid);
12. Had any of the following cervical pathology: high-grade cervical neoplasia, atypical glandular cells, atypical endocervical cells, atypical squamous cells favoring high grade. Patients with atypical squamous cells of undetermined significance and low-grade cervical neoplasia were allowed in the study if high-risk human papilloma virus testing was negative or if deoxyribonucleic acid testing for human papilloma virus 16 and 18 was negative;
13. Had any of the following clinical laboratory abnormalities at any screening visit:
- a. Hemoglobin <8.0 g per deciliter (patients with screening hemoglobin results <8 g per deciliter may have been prescribed iron supplements and had their hemoglobin levels retested prior to the baseline Day 1 visit);
 - b. Alanine aminotransferase (ALT) or aspartate aminotransferase (AST) >2.0 times the upper limit of normal (ULN), or bilirubin (total bilirubin) >1.5 times ULN on clinical laboratory testing at either the screening 1 or screening 2 visit (or >2.0 times ULN if secondary to Gilbert syndrome or pattern consistent with Gilbert syndrome);
 - c. Estimated glomerular filtration rate <60 ml/min/m² using the Modification of Diet and Renal Disease method;
 - d. Hypocalcemia (< lower limit of normal [LLN] or hypocalcemia (>ULN);
 - e. Hypophosphatemia (<LLN) or hyperphosphatemia (>ULN);
14. Had clinically significant cardiovascular disease, including the following:
- a. Prior history of myocardial infarction;
 - b. History of angina or significant coronary artery disease (i.e., ≥50% stenosis);
 - c. History of congestive heart failure;
 - d. History of clinically significant ventricular arrhythmias, such as ventricular tachycardia, ventricular fibrillation, or torsades de pointes, Mobitz II second degree or third degree heart block without permanent pacemaker in place or untreated supraventricular tachycardia (heart rate ≥120 beats per minute);
 - e. QT interval by the Fridericia correction (QTcF) of >470 msec on the screening visit or baseline Day 1 electrocardiogram;
 - f. Hypotension, as indicated by a systolic blood pressure <84 mmHg on two repeat measures at least 15 minutes apart, or treated ongoing symptomatic orthostatic hypotension with >20 mmHg decrease in systolic blood pressure one minute or more after assuming an upright

- position;
- g. Uncontrolled hypertension, as indicated by systolic blood pressure >160 mmHg or diastolic blood pressure >100 mmHg on two repeat measures at least 15 minutes apart at any screening visit or baseline Day 1 visit;
 - h. Bradycardia, as indicated by a heart rate <45 beats per minute on the screening electrocardiogram unless judged by the investigator to be due to physical fitness;
15. Had been a participant in an investigational drug or device study within the one month prior to the screening 1 visit;
 16. Had a history of clinically significant condition(s), including but not limited to the following:
 - a. Untreated thyroid dysfunction (patients with adequately treated hypothyroidism who were stable on medication were not excluded);
 - b. History of malignancy within the past five years or ongoing malignancy other than curatively treated nonmelanoma skin cancer or surgically cleared Stage 0 in situ melanoma;
 - c. Any current psychiatric disorder that would, in the opinion of the investigator or medical monitor, impair the ability of the patient to participate in the study or would impair interpretation of their data. Patients with major depression, post-traumatic stress disorder, based on the Diagnostic and Statistical Manual of Mental Disorders-5 criteria who had been unstable or not well controlled based on the investigator's or mental health professional's judgment or whose history or stability could not be ascertained, or whose psychiatric drug regimen had changed during the three months prior to screening or was expected to change during the study should not have been enrolled;
 - d. Had a systemic autoimmune disease (e.g., systemic lupus erythematosus, Sjogren syndrome, rheumatoid arthritis, polymyositis, systemic sclerosis, psoriasis, psoriatic arthritis, vasculitic syndromes, etc). Psoriasis not requiring or anticipated to require systemic therapy was permitted;
 17. Was pregnant or lactating, or intended to become pregnant during the study period through one month after the last dose of study drug or intended to donate ova during the study period or within two months after the last dose of study drug;
 18. Was using any prohibited medications;
 19. Had a contraindication or history of sensitivity to any of the study treatments or components thereof; or had a history of drug or other allergy that, in the opinion of the investigator or medical monitor, contraindicated study participation;
 20. Had a prior (within one year of screening 1 visit) or current history of drug or alcohol abuse disorder according the Diagnostic and Statistical Manual of Mental Disorders-5 (all patients had to be questioned about their drug and alcohol use and this should have been documented in the electronic case report form);
 21. Had participated in a previous clinical study that included the use of relugolix;
 22. Was an immediate family member, study site employee, or was in a dependent relationship with a study site employee who was involved in the conduct of the study (e.g., spouse, parent, child, or sibling);
 23. Was inappropriate for participation in this study because of conditions that may have interfered with interpretation of study results or prevent the patient from complying with study requirements, including contraception requirements, as determined by the investigator, subinvestigator, or medical monitor;
 24. Had received a blood transfusion within eight weeks prior to the screening 1 visit or during the screening period.

Study Objectives and End Points

The objectives and end points of the study are listed in the following table. All end points were prespecified in the statistical analysis plan, which was finalized prior to database lock and unblinding of the data. These analyses differ from what was originally included in the protocol, as described in the statistical analysis plan.

Table S2. Study Objectives and End Points.

Objective(s)	End Point(s)
Primary Efficacy	
To determine the benefit of relugolix 40 mg once a day co-administered with estradiol 1 mg and norethindrone acetate 0.5 mg compared with placebo for 24 weeks on heavy menstrual bleeding associated with uterine fibroids	Proportion of women in the relugolix combination therapy group versus the placebo group who achieve an MBL volume of <80 ml AND at least a 50% reduction from baseline menstrual blood loss volume over the last 35 days of treatment, as measured by the alkaline hematin method
Key Secondary Efficacy	
(Alpha-Protected for Hierarchical Hypothesis Testing — Relugolix Combination Therapy versus Placebo)	
Achievement of amenorrhea	Proportion of women who achieve amenorrhea over the last 35 days of treatment
Heavy menstrual bleeding associated with uterine fibroids	Percent change from baseline to week 24 in menstrual blood loss volume
Impact of uterine fibroids on symptoms, activities and health-related quality-of-life as measured by components of the UFS-QoL	Change from baseline to week 24 in the UFS-QoL bleeding and pelvic discomfort scale score, a sub-scale of the UFS-QoL Symptom Severity scale
Change in hemoglobin	Proportion of women with a hemoglobin ≤ 10.5 g per deciliter at baseline who achieve an increase of >2 g per deciliter from baseline at week 24
Pain associated with uterine fibroids	Proportion of patients with a maximum NRS score ≤ 1 during the 35 days before the last dose of study drug in the subset of women with a maximum NRS score ≥ 4 for pain associated with uterine fibroids during the 35 days prior to randomization
Uterine fibroid volume	Percent change from baseline to week 24 in volume of the largest uterine fibroid at baseline
Uterine volume	Percent change from baseline to week 24 in uterine volume

Objective(s)	End Point(s)
Other Secondary Efficacy (Not for Hierarchical Hypothesis Testing)*	
To determine the benefit of relugolix 40 mg once a day for 12 weeks followed by 12 weeks of relugolix 40 mg once a day co-administered with estradiol 1 mg and norethindrone acetate 0.5 mg compared with placebo for 24 weeks on heavy menstrual bleeding associated with uterine fibroids	Proportion of women in the delayed relugolix combination therapy group versus the placebo group who achieve a menstrual blood loss volume of <80 ml AND at least a 50% reduction from baseline menstrual blood loss volume over the last 35 days of treatment, as measured by the alkaline hematin method
Heavy menstrual bleeding associated with uterine fibroids	<ul style="list-style-type: none"> • Percent change from baseline to Weeks 4, 8, 12, 16, and 20 in menstrual blood loss volume • Change from baseline in menstrual blood loss volume by visit • Time to achieve an menstrual blood loss volume of <80 ml AND at least a 50% reduction from baseline menstrual blood loss volume as measured by the alkaline hematin method • Proportion of women in the relugolix combination therapy group versus the placebo group who achieve an menstrual blood loss volume of <80 ml AND at least a 50% reduction from baseline menstrual blood loss volume by visit
Achievement of amenorrhea	<ul style="list-style-type: none"> • Sustained amenorrhea rate by visit • Time to achieving sustained amenorrhea • Time to achieving amenorrhea
Change in hemoglobin	<ul style="list-style-type: none"> • Proportion of women who with a hemoglobin concentration below the lower limit of normal at baseline who achieved an increase of ≥ 1 g per deciliter from baseline at week 24 • Change from baseline to week 24 in hemoglobin for women with a hemoglobin concentration ≤ 10.5 g per deciliter at baseline
Impact of uterine fibroids on symptoms, activities, and health-related quality-of-life as measured by components of the UFS-QoL	<ul style="list-style-type: none"> • Change from baseline to week 24 in the UFS-QoL symptom severity scale score • Change from baseline to week 24 in the UFS-QoL activities scale score • Change from baseline to week 24 in the UFS-QoL revised activities scale score • Proportion of responders who achieved a meaningful increase of at least 20 points from baseline to week 24 in UFS-QoL revised activities scale score • Proportion of responders who achieved a meaningful reduction of at least 20 points from baseline to week 24 in UFS-QoL bleeding and pelvic discomfort scale score • Change from baseline to week 24 in the interference of uterine fibroids with physical activities based on UFS-QoL Question 11 • Change from baseline to week 24 in the interference of uterine fibroids with social activities based on UFS-QoL Question 20

Objective(s)	End Point(s)
	<ul style="list-style-type: none"> Change from baseline to week 24 in embarrassment caused by uterine fibroids based on UFS-QoL Question 29
Patient global assessment for function and symptoms as measured by the PGA for function and symptoms	<ul style="list-style-type: none"> Change in PGA for uterine fibroid-related function from baseline to week 24 Change in PGA for uterine fibroid symptoms from baseline to week 24 Proportion of patients achieving improvement from baseline in PGA for uterine fibroid symptoms from baseline to week 24 Proportion of patients achieving improvement from baseline in PGA for uterine fibroid related function from baseline to week 24
Impact of heavy menstrual bleeding on social, leisure, and physical activities as measured by the Menorrhagia Impact Questionnaire	<ul style="list-style-type: none"> Change from baseline to week 24 in the Menorrhagia Impact Questionnaire Score for physical activities Change from baseline to week 24 in the Menorrhagia Impact Questionnaire Score for social and leisure activities
Pain associated with uterine fibroids	Proportion of women who achieve a <i>maximum</i> NRS score for pain associated with uterine fibroids over the last 35 days of treatment that is at least a 30% reduction from baseline in the subset of women with a maximum pain score ≥ 4 during the 35 days prior to randomization
Safety	
To determine the safety of 24 weeks of relugolix 40 mg once a day co-administered with either 12 or 24 weeks of estradiol 1 mg and norethindrone acetate 0.5 mg in women with heavy menstrual bleeding associated with uterine fibroids compared with placebo for 24 weeks	Treatment-emergent adverse events (hereafter referred to as adverse events, change in vital signs (including weight), clinical laboratory tests, and electrocardiograms
To determine the percent change from baseline to week 12 in bone mineral density at the lumbar spine (L1 - L4) in the relugolix combination therapy group compared with the delayed relugolix combination therapy group in women with heavy menstrual bleeding associated with uterine fibroids	Percent change from baseline to week 12 in bone mineral density at the lumbar spine (L1 - L4) in the relugolix combination therapy group compared with the delayed relugolix combination therapy group, as assessed by dual-energy x-ray absorptiometry
To determine the change in bone mineral density of women with heavy menstrual bleeding associated with uterine fibroids treated with 24 weeks of relugolix 40 mg once a day co-administered with either 12 or 24 weeks of estradiol 1 mg and norethindrone acetate 0.5 mg compared with placebo for 24 weeks	Percent change from baseline to week 24 in bone mineral density at the lumbar spine (L1 - L4), total hip, and femoral neck as assessed by dual-energy x-ray absorptiometry

Objective(s)	End Point(s)
To determine the incidence of vasomotor symptoms with relugolix 40 mg once a day co-administered with either 12 or 24 weeks of estradiol 1 mg and norethindrone acetate 0.5 mg in women with heavy menstrual bleeding associated with uterine fibroids	Incidence of vasomotor symptoms
Pharmacokinetic and Pharmacodynamic	
To evaluate the pharmacokinetic and pharmacodynamic effects of 24 weeks of relugolix 40 mg once a day when co-administered with either 12 or 24 weeks of estradiol 1 mg and norethindrone acetate 0.5 mg	<ul style="list-style-type: none"> • Predose trough concentrations (C_{τ}) of relugolix, and NET and baseline-adjusted estradiol concentration • Absolute and changes from baseline to week 24 in predose concentrations of luteinizing hormone, follicle-stimulating hormone, estradiol, and progesterone
Exploratory	
To determine the benefit of 24 weeks of relugolix 40 mg once a day co-administered with either 12 or 24 weeks of estradiol 1 mg and norethindrone acetate 0.5 mg compared with placebo on patient-reported quality-of-life outcome measures (EQ-5D-5L)	Change from baseline to week 24 in the EQ-5D-5L scale

EQ-5D-5L denotes EuroQoL five-dimensions questionnaire (five-level version), NRS, numerical rating scale; PGA, patient global assessment, and UFS-QoL, Uterine Fibroid Symptom and Health-Related Quality of Life Questionnaire.

* The secondary end points below were assessed comparing relugolix Group A to placebo Group C inferentially; relugolix Group A to relugolix Group B and relugolix Group B to placebo Group C descriptively, unless otherwise specified.

Mixed-Effects Model for Imputing Missing Menstrual Blood Loss Volume Data

For the primary analysis, patients with missing menstrual blood loss volumes at week 24/end of treatment were identified per missing data handling rules. For imputing missing data for the primary analysis, a mixed-effects model approach was used, as the mixed-effects approach may better describe the effects of a hormonal treatment (such as suppression of the hypothalamic-pituitary-ovarian axis by GnRH antagonists).

Specifically, a mixed-effects model with repeated measures of menstrual blood loss volumes at multiple time points (Weeks 4, 8, 12, 16, 20 and 24) was fitted to predict percent change in menstrual blood loss volume from baseline (as a dependent variable) through the fixed-effects associated with covariates (i.e., stratification factors of baseline menstrual blood loss volume and geographic region, visit, treatment, and visit by treatment interaction) and random effects (from the individual patients). In this model, an unstructured variance-covariance matrix was assumed for each patient.

See sample SAS codes below for illustration where PCHG_MBL is percent change in menstrual blood loss volume from baseline as a dependent variable, PID is patient identification number, BMBL is a randomization stratification factor (baseline MBL <225 vs. ≥225), REGION is a randomization stratification factor (North America vs. Rest of World), TRT is treatment group (relugolix combination therapy or Placebo), VISIT is visit time point (4, 8, 12, 16, 20, and 24 weeks) and TRT*VISIT is the visit by treatment interaction. The specification of type=UN implements unstructured variance-covariance matrix for an individual patient with multiple measures of menstrual blood loss volumes.

```
proc mixed data=MBL_dataset method=REML covtest;

class PID BMBL REGION TRT VISIT;

model PCHG_MBL= BMBL REGION VISIT TRT VISIT*TRT/s outp=ufmi_mixed_p covb;

repeated VISIT /type=UN subject=PID r;

lsmeans TRT/diff;

ods output SolutionF=mixparms CovB=mixcovb;
```

Applying this model over the observed longitudinal menstrual blood loss volume data, the fixed-effects were estimated and relationship of percent change in menstrual blood loss volume from baseline with the covariates was characterized by the fitted model. From the fitted model, the percent change in menstrual blood loss volume (whether missing or not) was predicted for each patient at each visit and in a particular stratum. The imputed menstrual blood loss volume was obtained by first multiplying the imputed percent change with the individual patient's baseline menstrual blood loss volume to the difference, and then adding the baseline menstrual blood loss volume to the difference.

The main reason for using percent change in menstrual blood loss volume over reported menstrual blood loss volume as a dependent variable in the mixed-effects model is that the percent change is part of the derivation of the primary end point. Secondly, the percent change is a normalized value adjusted for the baseline value and less influenced by baseline menstrual blood loss volume, and therefore it is a better metric to describe the relationship of menstrual blood loss volume reduction with hormonal treatment and to impute the missing volumes in a more robust fashion.

Since the purpose of using a mixed-effects model is imputing the missing blood loss volumes identified at week 24/end of treatment, the predicted menstrual blood loss volumes at the corresponding week 24/end of treatment visit were used to determine responder status. For patients without the need for imputation, their responder status was derived according to the algorithms laid out in Table S3. This imputation approach is consistent with the definition of responder at week 24/end of treatment for the primary analysis.

Derivation of Responder Status at Week 24/End of Treatment and Missing Data Handling Rules

For the evaluation of primary end point, missing data handling rules were implemented for deriving responder status at week 24/end of treatment as described below. The following elements were checked: duration of treatment exposure; compliance with menstrual product collection against the daily electronic diary, as measured by their menstrual product return rate compliance with electronic diary entry, defined as the proportion of electronic diary entry days over the length (days) of menstrual product collection interval for week 24/end of treatment visit; and reasons for no menstrual product collection (as displayed in Table S3).

Patients with <4 weeks of treatment who withdrew from the study prematurely due to lack of efficacy or withdrew from the study prematurely to undergo surgical intervention for uterine fibroids were considered as non-responders.

All other patients had their responder status determined as follows:

- For patients with a menstrual product return rate of 100%, responder status was determined based on the observed menstrual blood loss volume;
- For patients who had incomplete menstrual product collection, with a menstrual product return rate of <100%, responder status was derived based on either imputed or observed menstrual blood loss volume;
 - Those with a menstrual blood loss volume ≥ 80 ml or $< 50\%$ reduction from baseline were considered as non-responders;
 - Those with a menstrual blood loss volume < 80 ml and $\geq 50\%$ reduction from baseline were imputed for partial or complete missing menstrual blood loss volume.
- For patients who did not return a menstrual product collection, responder status was determined depending on the reason reported on the Feminine Product Collection eCase Report Form:
 - If the reason was reported as Amenorrhea, the last 35 days of treatment were used to derive responder status:
 - If the week 24/end of treatment interval was 35 days, then she was considered as a responder;
 - If the week 24/end of treatment interval was < 35 days, the following supportive information was used to derive responder status:
 - If a patient reported amenorrhea at the visit prior to week 24/end of treatment, she was defined as a responder;
 - If a patient did not report amenorrhea at the visit prior to week 24/end of treatment, electronic diary data from the prior visit interval was reviewed to confirm whether the patient was amenorrheic for a total of 35 days.

- If the electronic diary from the previous interval confirmed amenorrhea, then the patient was considered as a responder;
 - Otherwise, menstrual blood loss volume was imputed.
- If the reason was Other and the specification described spotting or negligible bleeding, responder status was defined as follows:
 - The patient was considered as a responder if it was supported by the electronic diary data: the electronic diary entry rate must have exceeded 70% and the patient must have reported no more than 5 total days of bleeding with product use and no more than 3 consecutive bleeding with product use over the collection interval.
 - If the electronic diary entries did not confirm spotting or negligible bleeding, but the patient had at least 8 weeks of menstrual blood loss volume data prior to the week 24/end of treatment visit, her missing menstrual blood loss volume was imputed to determine responder status. Eight weeks of menstrual blood loss volume data represented a reasonable minimum length of observation to justify imputation of the remaining data in assessing the effects of hormonal therapy.
 - Otherwise if the patient had <8 weeks of menstrual blood loss volume data, she was considered as a non-responder;
- If the reason was any Other, the responder status was derived as follows:
 - If the patient had at least 8 weeks of menstrual blood loss volume data prior to the week 24/end of treatment visit, her missing menstrual blood loss volume was imputed and her responder status was based on the imputed menstrual blood loss volume.
 - If the patient had <8 weeks of menstrual blood loss volume data, she was considered as a non-responder.

Table S3. Derivation of Responder Status at Week 24/End of Treatment and Missing Data Handling Rules – for Primary Analysis.

Treatment Exposure	Menstrual Product Return Rate	Observed Menstrual Blood Loss Volume	Reason for No Menstrual Product Collection	Responder Status	
<4 weeks	N/A	N/A	N/A	Imputed as non-responder	
≥4 weeks	100% Menstrual Product Compliance	N/A	N/A	Based on the observed menstrual blood loss volume	
	<100% Menstrual Product Compliance	Menstrual blood loss volume ≥80 ml or <50% reduction from baseline	N/A	Imputed as non-responder based on the observed menstrual blood loss volume	
		Menstrual blood loss volume <80 ml and ≥50% reduction from baseline	N/A	Based on the imputed menstrual blood loss volume	
	No Menstrual Product Collection	N/A	N/A	Reported “Amenorrhea”	Imputed as responder
				Reported “Spotting or negligible bleeding” and confirmed by eDiary*	Imputed as responder
				Reported “Spotting or negligible bleeding” although not confirmed by eDiary or any other reason, had at least 8 weeks of menstrual blood loss volume data	Based on the imputed menstrual blood loss volume
				The entries in the eDiary did not verify “Spotting or negligible bleeding” or any other reason and if had <8 weeks of menstrual blood loss volume data	Imputed as non-responders

eDiary denotes electronic diary, and N/A not available.


* Defined as those patients who meet the following criteria: eDiary entry rate >70% and no more than 3 consecutive days and no more than 5 days of bleeding/spotting and use of menstrual product reported on the eDiary over the week 24/end of treatment visit window.

Key Secondary Efficacy End Points with Alpha-Protection

For testing whether relugolix combination therapy (Group A) was statistically significantly superior to placebo (Group C) for the primary efficacy end point as well as the seven key secondary end points listed below, a gate-keeping mixed sequence testing procedure was applied to maintain the family-wise type I error rate. Under this testing procedure, the primary end point was tested first at a 2-sided 0.05 significance level. If the P-value for primary end point was <0.05 , the seven key end points listed below were tested sequentially in the order depicted in Figure S1 and Figure S2.


For the relugolix combination therapy group to be considered statistically superior to the placebo group on a secondary end point, the two-sided P-value must be <0.05 for that secondary end point and for all higher-ranking secondary end points, as well as for the primary end point. If the two-sided P-value was <0.05 for the fourth end point (proportion of women with a hemoglobin ≤ 10.5 g per deciliter at baseline who achieve an increase of >2 g per deciliter from baseline at week 24 for LIBERTY 1; proportion of women who achieve a maximum NRS score ≤ 1 for uterine fibroid-associated pain over the last 35 days of treatment in the subset of women with a maximum pain score ≥ 4 during the 35 days prior to randomization for LIBERTY 2), the remaining three end points (the fifth, sixth, or seventh) were tested using the Hochberg step-up procedure. The fourth end point tested for LIBERTY 2 (Figure S2) was changed (i.e., reordered relative to LIBERTY 1) after results became available from LIBERTY 1 (Figure S1) but before data unblinding of LIBERTY 2.

Figure S1. Mixed Sequence Testing Procedure for Primary and Key Secondary End Points in LIBERTY 1.



BPD denotes Bleeding and Pelvic Discomfort, EP end point, Hgb hemoglobin, max maximum, MBL menstrual blood loss, M-vol myoma volume, NRS Numerical Rating Scale, PE primary end point, Prop proportion, UFS-QoL BPD Uterine Fibroid Symptom and Health-Related Quality of Life Bleeding and Pelvic Discomfort, and U-vol uterine volume.

Figure S2. Mixed Sequence Testing Procedure for Primary and Key Secondary End Points in LIBERTY 2.



From the Hochberg procedure, the P-values were calculated for the three end points (5, 6, and 7 for LIBERTY 1 and 4, 6, and 7 for LIBERTY 2) and ranked from the smallest to the largest. The end point corresponding to the largest P-value gets tested first. If the P-value was < 0.05 , then no further testing occurred, and it was concluded that all three end points are positive. Otherwise, the end point corresponding to the second largest P-value was tested. If the P-value was < 0.025 , then no further testing occurred, and it was concluded that the end points corresponding to the middle and smallest P-values are positive. Otherwise, the end point with the smallest P-value was tested. If the P-value was < 0.0167 , no further testing occurred, and it was concluded that only the end point with the smallest P-value was positive. Otherwise, all three end points did not pass the statistical significance criterion at 0.05 level.

The seven key secondary efficacy end points are numbered as follows:

1. Proportion of women who achieve amenorrhea over the last 35 days of treatment;
2. Percent change from baseline to week 24 in MBL volume;
3. Change from baseline to week 24 in Bleeding and Pelvic Discomfort Scale score as measured by the UFS-QoL Symptom Severity Scale (Q1, Q2, Q5);

4. Proportion of women with a hemoglobin \leq 10.5 g per deciliter at baseline who achieve an increase of >2 g per deciliter from baseline at week 24
5. Proportion of women who achieve a maximum NRS score \leq 1 for uterine fibroid associated pain over the last 35 days of treatment in the subset of women with a maximum pain score ≥ 4 during the 35 days prior to randomization;
6. Percent change from baseline to week 24 in uterine fibroid volume;
7. Percent change from baseline to week 24 in uterine volume.

For key secondary efficacy end points (1, 4, and 5) that are evaluating proportions, treatment comparisons were performed using a stratified Cochran-Mantel-Haenszel test with the randomization stratification factors as strata. Point estimates and 2-sided 95% CIs for treatment differences in proportions were provided.

For key secondary end point 4, an increase in hemoglobin of 2g per deciliter was considered clinically meaningful, because it corresponds to approximately the same increase as that expected after a transfusion of ~ 2 units of packed red blood cells (Man, 2016; Bachowski, 2017).

For deriving the key secondary end point 5 (proportion of women who achieve a maximum NRS score ≤ 1 for uterine fibroid-associated pain over the last 35 days of treatment in the subset of women with a maximum pain score ≥ 4 during the 35 days prior to randomization), the patient subset and week 24/end of treatment maximum value are determined as follows.

Because patients were asked to begin electronic diary entries after returning the first collection of menstrual products, the number of electronic diary entries made during screening varies with the duration of screening for each patient. Some patients required only one collection to be randomized, whereas others required as many as four collections to confirm eligibility.

Once the qualifying menstruation was completed and the patient qualified for randomization based upon resulting MBL volume(s), the recording of patient's NRS scores for screening phase was ended and the number of pain score days at baseline could be as short as 7 days or as long as 70 days prior to randomization. If a patient met the subset definition (maximum NRS score ≥ 4 at baseline) over a portion of the screening days (e.g., 7–70 days), she also met the subset definition on the entire 35 days interval.

Since the maximum NRS value was used to determine inclusion into the subset rather than an average NRS value, the variable number of days for inclusion of patients has no major impact on determining patient subset. To ensure robust estimate of response, the minimum number of non-missing daily pain scores required to calculate the maximum score at week 24/end of treatment was at least 28 days (80% of the last 35 days of treatment) of pain scores recorded in the e-Diary entry.

The primary analysis of key secondary end point 5 was analyzed for the subset of women who had a maximum pain score ≥ 4 during the 35 days prior to randomization and who had at least 28 days (80% of the last 35 days of treatment) of pain scores recorded in the e-Diary at week 24/end of treatment. In addition, a sensitivity analysis was conducted on the subset of women who had a maximum pain score ≥ 4 during the 35 days prior to randomization without restricting number of days of pain scores recorded in the e-Diary.

The analysis for end point 5 (proportion of women who achieve a maximum NRS score ≤ 1 for uterine fibroid-associated pain over the last 35 days of treatment in the subset of women with a maximum pain score ≥ 4 during the 35 days prior to randomization) was also performed using NRS scores reported on electronic diary during menstrual and non-menstrual days.

For key secondary efficacy end points (6 and 7) evaluating percent change from baseline in uterine fibroid volume and uterine volume that are measured only at week 24, an analysis of covariance (ANCOVA) model was used to assess treatment effect with treatment, randomization stratification factors and baseline value as covariates.

For key secondary efficacy end points (2 and 3) evaluating the change (absolute or % change) from baseline to week 24 by ultrasound, treatment comparisons were performed using a mixed model repeated measures approach with treatment, visit, randomization stratification factors and treatment by visit interactions included as fixed effects and random effects (from the individual patients). In this model, an unstructured variance-covariance matrix was assumed for each patient.

Patient-Reported Outcomes

Overview

The justification for evaluating patient-reported outcomes in women with uterine fibroids is the clinical importance of symptomatology. Specifically, the condition may manifest as pain, symptoms related to bleeding and to fibroid bulk, and/or attendant functional, social, and emotional effects that adversely affect quality of life.

In the phase 2 study Japanese study in uterine fibroids, a dose-dependent trend in improvement of pain associated with uterine fibroids was recognized that warranted further investigation in phase 3. As a consequence, an important goal of the pivotal relugolix combination therapy and relugolix monotherapy studies was to more fully characterize the impact of relugolix on pain and other patient-reported outcomes in women with uterine fibroids.

In the relugolix combination therapy LIBERTY studies, a key secondary objective was to describe the effect of relugolix in combination with estradiol and norethindrone acetate compared with placebo on pain associated with uterine fibroids. Pain associated with uterine fibroids was measured daily (i.e., during menstrual as well as non-menstrual days) using the well-recognized and validated numerical rating scale (NRS) by electronic diary. Another key secondary end point was distress due to heavy menstrual bleeding, passing blood clots, and pelvic tightness symptoms (as measured by a validated instrument, the Bleeding and Pelvic Discomfort [BPD] scale [Questions 1, 2, and 5 from Uterine Fibroid Symptoms-Quality of Life [UFS-QoL]).

In the phase 3 Japanese relugolix monotherapy program, a separate study of relugolix in women with moderate to severe pain associated with uterine fibroids was conducted in which pain was the primary end point using the NRS instrument (Osuga, 2019).

Specific Assessments and Instruments

1. Pain: Numerical Rating Scale (NRS)

Selection of Instrument and Method of Assessment

Pain is the second most frequent and debilitating symptom for women with uterine fibroids (David et al. 2016; Foth et al. 2017, Monleon et al. 2018) and therefore was identified as an important end point to measure in the relugolix development program.

Women's pain experience is individual and covers a range of pain symptoms, with dysmenorrhea and pelvic pain frequently mentioned (Donnez et al. 2016). Since pain experience is patient specific, and patients spontaneously report different types of pain (Deal et al. 2011), a general measure of pain associated with uterine fibroids was deemed to be best suited and most appropriate for assessment in the relugolix clinical program.

The pain NRS is a validated, single-item, self-reported measure, which asks respondents to rank their pain on an 11-point scale (Ameade and Mohammed 2016). In chronic pain populations, patients have been found to prefer the NRS over other measures of pain intensity due to the NRS's comprehensibility and ease of completion (Hawker et al. 2011). The NRS measures have been described as less abstract and easier to understand than the visual analogue scale (VAS) (Dworkin et al. 2005). In a summary of studies using different pain scales, the NRS was found to have high compliance rates (Hjermstad et al. 2011). The NRS has been found to be highly correlated to the VAS in patients with rheumatic and other

chronic pain conditions (pain >6 months): correlations range from 0.79 to 0.95, suggesting that both the NRS and VAS are measuring the same concept (Kahl and Cleland 2005; Hawker et al. 2011).

In the relugolix clinical program, women were asked to document in a patient electronic diary the worst pain associated with their uterine fibroids that they had experienced during the last 24 hours on a scale from 0 to 10, with 0 indicating “no pain” and 10 indicating “pain as bad as you can imagine.” Recording of “worst pain” helps mitigate the confounding issue of timing of the reporting of the NRS versus timing of analgesic administration, the latter of which would be expected to lower the pain score.

End Points

In the pivotal relugolix combination therapy LIBERTY studies, the proportion of women who achieved a maximum NRS score for pain associated with uterine fibroids over the last 35 days of treatment that was ≤ 1 (minimal to no pain) was a key secondary end point in the subset of women with a maximum pain score of ≥ 4 (moderate to severe pain) during the time period prior to randomization and who completed the daily electronic diary during the 35 days prior to randomization (pain-evaluable population).

Similarly, in the relugolix Japanese study, which enrolled only women with a maximum pain score of ≥ 4 (moderate to severe pain) at baseline, the proportion of women who achieved a maximum NRS score for pain associated with uterine fibroids of ≤ 1 (minimal to no pain) over the last 28 days of treatment was reported as the primary end point.

NRS Thresholds for Analyses

The threshold of a maximum pain score of ≤ 1 was chosen for the phase 3 studies because patient reports of minimal to no pain would represent the maximal improvement possible. Analgesic use was analyzed for the pain-evaluable population as supportive data to the primary analysis of the proportion of patients meeting the NRS threshold.

The pivotal studies with relugolix combination therapy also assessed as a supportive secondary end point, the percentage of patients with a 30% reduction from baseline in NRS score. Traditionally, pain reductions from baseline of at least 30% have been used as a threshold of clinically meaningful change in chronic pain clinical studies (Dworkin et al. 2005).

2. Bleeding and Pelvic Discomfort: BPD Scale

A key secondary end point in the pivotal relugolix combination therapy studies was the change from baseline to week 24 in the UFS-QoL BPD scale score, with a range of possible scores from 0 to 100, where higher scores are indicative of greater distress and lower scores are indicative of less distress.

The UFS-QoL BPD scale was derived and validated by Myovant using exploratory and confirmatory factor analysis. It assesses distress experienced by patients due to three symptoms associated with uterine fibroids that are common to most patients (i.e., heavy bleeding during the menstrual period [Question 1], passing blood clots during the menstrual period [Question 2], and feeling tightness or pressure in the pelvic area [Question 5]) (Spies et al. 2002). Questions 1 and 2 are closely related to the primary end point, they provide the patient perspective in terms of distress due to heavy bleeding and passing blood clots, while the primary end point objectively measures blood loss volume. Question 5 is the expected patient symptomatology that is related to the uterine volume end point.

The meaningful change threshold for the BPD was derived via anchor-based analyses using pooled blinded data from the first third of patients enrolled in one of the two LIBERTY studies, who completed


week 24 visits and had PGA scores for symptoms at baseline and week 24. Findings were supported by the results from the qualitative exit interview study, in which the patients' perception of what constitutes a meaningful change on the BPD scale was assessed. The meaningful change threshold was set at 20 points, based on the totality of data from anchor-based analyses from blinded phase 3 data, supported by cumulative distribution function and probability density function curves as well as the results from the exit interview study. As illustrated in the cumulative distribution function curves (Supplementary Methods), setting the meaningful change threshold at 20-point improvement on the BPD scale would yield a percent of misclassified responders for the "no change" group of 33% and a percent of correctly classified responders of 60% for the "1 category improvement" group.

The BPD scale was only assessed as an end point in the relugolix combination therapy LIBERTY studies and not in the relugolix monotherapy program.

Description of development and validation of the BPD instrument has been presented (Li et al. 2019) and is under preparation as a separate manuscript.

Supplementary Figures


Figure S3. Study Design.



E2 denotes estradiol, NETA norethindrone acetate, and QD once a day.


* E2 1 mg, NETA 0.5 mg.

Figure S4. Patient Disposition.



Patients not meeting eligibility criteria due to lack of confirmation of uterine fibroids or heavy menstrual bleeding: LIBERTY 1, 971 (63.4%); LIBERTY 2, 1202 (57.4%).


Figure S5. Subgroup Analyses for Primary End Point.*



* Primary end point is the proportion of responders with menstrual blood loss volume < 80 mL and ≥ 50% reduction from baseline over the last 35 days of treatment. All variables are at baseline. Unless otherwise noted, odds ratios are based on logistic regression with treatment group, baseline menstrual blood loss volume, and geographic region (North America, Rest of World) as covariates. CI denotes confidence interval and NE not estimable. 95% CIs are not adjusted for multiplicity and thus should not be used to infer definitive treatment effects.



† Odds ratio based on logistic regression with treatment group as the only covariate due to smaller sample size.

Figure S6. Secondary Efficacy End Points: A) Reduction in Menstrual Blood Loss, B) Relief of Pain Associated With Uterine Fibroids, and C) Reduction in Bleeding and Pelvic Discomfort.



In A, least squares means and P value for test of difference of relugolix combination therapy minus placebo was based on a mixed-effects model with baseline menstrual blood loss, region, treatment, visit, and treatment by visit as fixed effects; lines are staggered for visibility. In B, results are analyzed for the patient subgroup with moderate/severe pain (Numerical Rating Scale score ≥ 4) associated with uterine fibroids during the 35 days prior to randomization and at least 28 days of electronic daily diary entries during the last 35 days of treatment. In C, the Bleeding and Pelvic Discomfort transformed score ranges from 0 to 100, with higher scores indicating greater symptom severity. CI denotes confidence interval.

*Data shown for LIBERTY 2 are from a sensitivity analysis excluding a week-4 menstrual blood loss volume of 2710 mL for one woman in the relugolix combination therapy arm (menstrual blood loss data from this patient were included at all other time points). The week-4 least squares mean menstrual blood loss change from baseline in the pre-specified analysis with this outlying week-4 data point included was 36.1% (95% CI 57.2%, 15.0%; nominal P=0.055 vs. placebo) and in the sensitivity analysis with this data point excluded was 52.1% (95% CI: 63.0%, 41.2%, nominal P<0.001 vs. placebo).



Supplementary Tables

Table S4. Additional Baseline Characteristics.

	LIBERTY 1			LIBERTY 2		
	Placebo (N = 127)	Relugolix Combination Therapy (N = 128)	Delayed Relugolix Combination Therapy (N = 132)	Placebo (N = 129)	Relugolix Combination Therapy (N = 125)	Delayed Relugolix Combination Therapy (N = 127)
Geographic Region – no. (%)						
North America	98 (77.2)	98 (76.6)	101 (76.5)	96 (74.4)	93 (74.4)	94 (74.0)
Rest of world	29 (22.8)	30 (23.4)	31 (23.5)	33 (25.6)	32 (25.6)	33 (26.0)
Hypertension – no. (%)	23 (18.1)	33 (25.8)	21 (15.9)	24 (18.6)	23 (18.3)	29 (23.0)

Table S5. Other Secondary End Points.

End Point Statistics	LIBERTY 1			LIBERTY 2		
	Placebo (N = 127)	Relugolix Combination Therapy (N = 128)	Delayed Relugolix Combination Therapy (N = 132)	Placebo (N = 129)	Relugolix Combination Therapy (N = 125)	Delayed Relugolix Combination Therapy (N = 127)
Percent change from baseline to week 24 in hemoglobin concentrations for women with hemoglobin ≤ 10.5 g per deciliter at baseline						
LS mean \pm SE	10.0 \pm 3.5	20.8 \pm 3.1	24.6 \pm 3.0	4.3 \pm 2.7	24.3 \pm 3.0	29.4 \pm 3.0
95% CI	3.0, 17.0	14.7, 26.8	18.6, 30.5	-1.1, 9.6	18.3, 30.3	23.5, 35.3
Difference of LS mean \pm SE*		10.8 \pm 4.2	14.6 \pm 4.2		20.0 \pm 3.9	25.1 \pm 3.9
95% CI		2.5, 19.2	6.4, 22.9		12.3, 27.8	17.4, 32.9
Patients with a hemoglobin concentration below the lower limit of normal† who achieved an increase of ≥ 1 g per deciliter from baseline to week 24						
Patients with hemoglobin below lower limit of normal at baseline – no. (%)	67 (52.8)	72 (56.3)	83 (62.9)	81 (62.79)	69 (55.2)	75 (59.1)
no. (%)	17 (25.4)	34 (47.2)	46 (55.4)	18 (22.2)	35 (50.7)	44 (58.7)
95% CI – %‡	15.53, 37.49	35.33, 59.35	44.10, 66.34	13.73, 32.83	38.41, 62.98	46.70, 69.92
Difference from placebo (95% CI) – %§		21.85 (6.31, 37.39)	30.05 (15.12, 44.98)		28.50 (13.63, 43.37)	36.44 (22.09, 50.80)
Responders with a meaningful reduction of ≥ 20 points from baseline to week 24 in BPD scale score						
no. (%)	35 (27.6)	79 (61.7)	83 (62.9)	37 (28.7)	79 (63.2)	69 (54.3)
95% CI – %‡	20.01, 36.19	52.72, 70.17	54.04, 71.12	21.07, 37.30	54.11, 71.65	45.26, 63.19
Difference from placebo (95% CI) – %§		34.16 (22.70, 45.62)	35.32 (23.99, 46.65)		34.52 (23.01, 46.02)	25.65 (13.99, 37.31)

BPD denotes Bleeding and Pelvic Discomfort, CI confidence interval, LS least squares, and SE standard error.

* Based on mixed-effects model with treatment, visit, region, baseline menstrual blood loss and treatment-by-visit interaction included as fixed effects. The multiple visits for each patient were the repeated measures as a random effect within each patient and an unstructured covariance.

† Lower limit of normal is hemoglobin < 11.6 g per deciliter.

‡ Based on exact binomial 95% CI (Clopper-Pearson).

§ Difference is Relugolix combination therapy or Delayed relugolix combination therapy minus Placebo. 95% CI for difference is based on the normal approximation not adjusted for multiplicity, and thus should not be used to infer definitive treatment effects.

Table S6. Serious Adverse Events.*

Serious adverse events – no. (%)	LIBERTY 1			LIBERTY 2		
	Placebo (N = 127)	Relugolix Combination Therapy (N = 128)	Delayed Relugolix Combination Therapy (N = 132)	Placebo (N = 129)	Relugolix Combination Therapy (N = 126)	Delayed Relugolix Combination Therapy (N = 126)
At least one	2 (1.6)	7 (5.5)	3 (2.3)	4 (3.1)	1 (0.8)	2 (1.6)
Ankle fracture	0	1 (0.8)	1 (0.8)	0	0	0
Avulsion fracture	0	1 (0.8)	0	0	0	0
Hematemesis	0	1 (0.8)	0	0	0	0
Hypothyroidism	0	1 (0.8)	0	0	0	0
Menorrhagia	0	1 (0.8)	0	0	0	0
Pelvic pain	0	1 (0.8)	0	0	0	0
Rhabdomyolysis	0	1 (0.8)	0	0	0	0
Uterine leiomyoma	0	1 (0.8)	0	0	0	0
Uterine myoma expulsion	0	1 (0.8)	0	0	0	0
Vitreous detachment	0	1 (0.8)	0	0	0	0
Acute psychosis	1 (0.8)	0	0	0	0	0
Appendicitis	0	0	1 (0.8)	0	0	0
Panic attack	0	0	1 (0.8)	0	0	0
Pneumonia	1 (0.8)	0	0	0	0	0
Cholecystitis	0	0	0	0	1 (0.8)	0
Anemia	0	0	0	1 (0.8)	0	0
Cholecystitis acute	0	0	0	0	0	1 (0.8)
Intervertebral disc degeneration	0	0	0	0	0	1 (0.8)
Intervertebral disc protrusion	0	0	0	0	0	1 (0.8)
Necrotising fasciitis	0	0	0	1 (0.8)	0	0
Radius fracture	0	0	0	1 (0.8)	0	0
Road traffic accident	0	0	0	1 (0.8)	0	0
Syncope	0	0	0	1 (0.8)	0	0

* Adverse events were coded using the *Medical Dictionary for Regulatory Activities* and severity of adverse events was evaluated by the investigator based on the National Cancer Institute's *Common Terminology for Adverse Events* (version 5.0).

Table S7. Percent Change from Baseline in Bone Mineral Density at Lumbar Spine and Total Hip in LIBERTY 1 and LIBERTY 2.

Location Time point Statistic	LIBERTY 1			LIBERTY 2		
	Placebo (N = 127)	Relugolix combination therapy (N = 128)	Delayed relugolix combination therapy (N = 132)	Placebo (N = 129)	Relugolix combination therapy (N = 126)	Delayed relugolix combination therapy (N = 126)
Lumbar spine						
Week 12						
no.	103	101	103	104	103	95
LS mean percent change from baseline (95% CI)	0.20 (-0.36, 0.76)	-0.47 (-1.04, 0.10)	-2.00 (-2.56, -1.43)	0.51 (-0.01, 1.03)	-0.82 (-1.35, -0.29)	-1.92 (-2.46, -1.37)
Week 24						
no.	102	100	100	95	95	94
LS mean percent change from baseline (95% CI)	0.05 (-0.52, 0.62)	-0.36 (-0.93, 0.22)	-1.82 (-2.39, -1.25)	0.32 (-0.26, 0.89)	-0.13 (-0.71, 0.46)	-2.12 (-2.71, -1.53)
Total hip						
Week 12						
no.	104	102	100	102	104	93
LS mean percent change from baseline (95% CI)	0.41 (-0.03, 0.85)	0.01 (-0.45, 0.46)	-0.95 (-1.40, -0.50)	-0.16 (-0.56, 0.24)	0.05 (-0.35, 0.45)	-1.07 (-1.48, -0.65)
Week 24						
no.	103	100	98	95	98	92
LS mean percent change from baseline (95% CI)	0.55 (0.08, 1.02)	0.02 (-0.46, 0.51)	-1.04 (-1.52, -0.56)	-0.04 (-0.48, 0.39)	-0.17 (-0.61, 0.26)	-1.16 (-1.60, -0.71)

CI denotes confidence interval and LS least squares.

Table S8. Summary of Patients With Liver Function Tests Meeting Predefined Limits of Change.

Patients – no. (%)	LIBERTY 1			LIBERTY 2		
	Placebo (N = 127)	Relugolix Combination Therapy (N = 128)	Delayed Relugolix Combination Therapy (N = 132)	Placebo (N = 129)	Relugolix Combination Therapy (N = 126)	Delayed Relugolix Combination Therapy (N = 126)
ALT or AST $\geq 3 \times$ ULN and $< 5 \times$ ULN	1 (0.8)	1 (0.8)	1 (0.8)	0	0	1 (0.8)
ALT or AST $\geq 5 \times$ ULN and $< 10 \times$ ULN	0	0	1 (0.8)	0	0	0
ALT or AST $\geq 10 \times$ ULN and $< 20 \times$ ULN	0	0	0	0	0	0
ALT or AST $\geq 20 \times$ ULN	0	0	0	0	0	0
BILI $\geq 2 \times$ ULN	0	0	0	0	0	0
ALT or AST $\geq 3 \times$ ULN and BILI $\geq 2 \times$ ULN	0	0	0	0	0	0

ALT denotes alanine aminotransferase, AST aspartate aminotransferase, BILI total bilirubin, and ULN upper limit of normal.

Table S9. Summary of Change from Baseline in Serum Lipids.

Parameter Statistics	LIBERTY 1			LIBERTY 2		
	Placebo (N = 127)	Relugolix Combination Therapy (N = 128)	Delayed Relugolix Combination Therapy (N = 132)	Placebo (N = 129)	Relugolix Combination Therapy (N = 126)	Delayed Relugolix Combination Therapy (N = 126)
Cholesterol – mg/deciliter						
Change from baseline at week 24						
no.	97	92	98	92	91	89
Mean ±SD	-1.9±25.3	-2.1±22.1	2.0±25.1	-0.8±20.8	0.8±31.6	5.5±25.5
Maximum increase from baseline during treatment						
no.	52	46	61	44	51	63
Mean ±SD	14.8±14.1	16.4±12.5	18.1±14.8	17.7±11.9	22.7±22.5	18.2±16.9
LDL cholesterol – mg/deciliter						
Change from baseline at week 24						
no.	97	92	96	92	90	89
Mean ±SD	-2.8±21.2	0.3±18.2	0.9±20.8	-0.7±17.4	2.6±28.3	4.6±21.9
Maximum increase from baseline during treatment						
no.	54	53	53	49	57	57
Mean ±SD	11.8±11.9	12.2±11.1	15.8±11.4	13.0±11.1	17.9±21.6	17.3±15.2
HDL cholesterol – mg/deciliter						
Change from baseline at week 24						
no.	97	92	98	92	91	89
Mean ±SD	0.1±9.7	-1.2±8.4	0.2±7.9	1.8±7.3	-1.0±7.9	0.3±7.0
Maximum decrease from baseline during treatment						
no.	50	58	54	47	60	44
Mean ±SD	-7.2±7.5	-6.3±5.2	-5.5±5.3	-4.6±3.8	-6.4±5.4	-4.5±3.6

Parameter Statistics	LIBERTY 1			LIBERTY 2		
	Placebo (N = 127)	Relugolix Combination Therapy (N = 128)	Delayed Relugolix Combination Therapy (N = 132)	Placebo (N = 129)	Relugolix Combination Therapy (N = 126)	Delayed Relugolix Combination Therapy (N = 126)
Triglycerides – mg/deciliter						
Change from baseline at week 24						
no.	97	92	98	92	91	89
Mean ±SD	5.2±44.8	-4.0±45.9	4.8±49.3	-9.1±50.0	-4.7±58.0	3.0±44.7
Maximum decrease from baseline during treatment						
no.	58	45	63	43	44	55
Mean ±SD	29.3±38.9	31.0±29.2	31.6±32.8	28.6±26.4	40.0±40.0	33.1±33.5

HDL denotes high-density lipoprotein, LDL low-density lipoprotein, and SD standard deviation.

References

- Ameade EPK, Mohammed BS. Menstrual pain assessment: comparing verbal rating scale (VRS) with numerical rating scales (NRS) as pain measurement tools. *Int J Womens Health Wellness*. 2016;2(1):017.
- David M, Pitz CM, Mihaylova A, Siedentopf F. Myoma-associated pain frequency and intensity: a retrospective evaluation of 1548 myoma patients. *Eur J Obstet Gynecol Reprod Biol*. 2016;199:137-140.
- Deal LS, Williams VS, Fehnel SE. Development of an electronic daily uterine fibroid symptom diary. *Patient*. 2011;4(1):31-44.
- Donnez J, Dolmans MM. Uterine fibroid management: from the present to the future. *Hum Reprod Update*. 2016;22(6):665-686.
- Dworkin RH, Turk DC, Farrar JT, Haythornthwaite JA, Jensen MP, Katz NP, et al. Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain*. 2005;113(1-2):9-19.
- Foth D, Rohl FW, Friedrich C, Tylkoski H, Rabe T, Romer T, et al. Symptoms of uterine myomas: data of an epidemiological study in Germany. *Arch Gynecol Obstet*. 2017;295(2):415-426.
- Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res (Hoboken)*. 2011;63 Suppl 11:S240-252.
- Hjermstad MJ, Fayers PM, Haugen DF, Caraceni A, Hanks GW, Loge JH, et al. Studies comparing numerical rating scales, verbal rating scales, and visual analogue scales for assessment of pain intensity in adults: a systematic literature review. *J Pain Symptom Manage*. 2011;41(6):1073-1093.
- Kahl C, Cleland JA. Visual analogue scale, numeric pain rating scale and the McGill Pain Questionnaire: an overview of psychometric properties. *Physical Therapy Reviews*. 2005;10(2):123-128.
- Li J, Kang JB, Hunsche E, Hudgens S. Measuring patient-reported outcomes in women with heavy menstrual bleeding associated with uterine fibroids: the Bleeding and Pelvic Discomfort scale. Annual scientific sessions of the American Society for Reproductive Medicine, Philadelphia, PA, October 12–16, 2019, P-595. *Fertil Steril* 2019;112 (3 Suppl):e344.
- Man L, Tahhan HR. Body surface area: a predictor of response to red blood cell transfusion. *J Blood Med*. 2016;7:199-204.
- Monleon J, Canete ML, Caballero V, Del Campo M, Domenech A, Losada MA, et al. Epidemiology of uterine myomas and clinical practice in Spain: An observational study. *Eur J Obstet Gynecol Reprod Biol*. 2018;226:59-65.
- Spies JB, Coyne K, Guaou Guaou N, Boyle D, Skyrnarz-Murphy K, Gonzalves SM. The UFS-QOL, a new disease-specific symptom and health-related quality of life questionnaire for leiomyomata. *Obstet Gynecol*. 2002;99(2):290-300.